

10/525783

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
12 May 2005 (12.05.2005)

PCT

(10) International Publication Number
WO 2005/042538 A1

(51) International Patent Classification⁷: **C07D 491/04, A61K 31/438, A61P 25/00**

(21) International Application Number:
PCT/GB2004/004484

(22) International Filing Date: 21 October 2004 (21.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/512,893 21 October 2003 (21.10.2003) US

(71) Applicant (for all designated States except MG, US): AS-TRAZENECA AB [SE/SE]; S-151 85 Södertälje (SE).

(71) Applicant (for MG only): ASTRAZENECA UK LIMITED [GB/GB]; 15 Stanhope Gate, London, Greater London W1K 1LN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): PHILLIPS, Eifion [GB/US]; AstraZeneca R & D Wilmington, 1800 Concord Pike, Wilmington, DE 19850-5437 (US).

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; AstraZeneca AB, S-151 85 Södertälje (SE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.


(54) Title: SPIROFUROPYRIDINE ARYL DERIVATIVES



(I)



(II)



(III)

(57) Abstract: Compounds of formula (I); and pharmaceutically-acceptable salts thereof, wherein: Ar is a moiety of formula (II) or (III); and A, B, and R¹ are as defined in the specification, compositions containing such compounds and the use of such compounds and compositions for use in therapy.

WO 2005/042538 A1

SPIROFUOPYRIDINE ARYL DERIVATIVES

TECHNICAL FIELD

This invention relates to novel spirofuopyridine aryl ligands for nicotinic
5 acetylcholine receptors and their use in therapy.

BACKGROUND OF THE INVENTION


The use of compounds which bind to nicotinic acetylcholine receptors for the
treatment of a range of disorders involving reduced cholinergic function such as Alzheimer's
10 disease, cognitive or attention disorders, anxiety, depression, smoking cessation,
neuroprotection, schizophrenia, analgesia, Tourette's syndrome, and Parkinson's disease is
discussed in: McDonald *et al.*, (1995) "Nicotinic Acetylcholine Receptors: Molecular
Biology, Chemistry and Pharmacology", Chapter 5 in Annual Reports in Medicinal
Chemistry, vol. 30, pp. 41-50, Academic Press Inc., San Diego, CA; Williams *et al.*, (1994)
15 "Neuronal Nicotinic Acetylcholine Receptors," Drug News & Perspectives, vol. 7, pp. 205-
223; Holladay *et al.* (1997) *J. Med. Chem.* 40(26), 4169-4194; Arneric and Brioni (Eds.)
(1998) "Neuronal Nicotinic Receptors: Pharmacology and Therapeutic Opportunities", John
Wiley & Sons, New York; Levin (Ed.) (2001) "Nicotinic Receptors in the Nervous System"
CRC Press.

20 Compounds which bind to nicotinic acetylcholine receptors and particularly those that
bind to alpha-7 nicotinic acetylcholine receptors are useful for the treatment of a range of
disorders involving reduced cholinergic function such as Alzheimer's disease, cognitive or
attention disorders, anxiety, depression, schizophrenia, analgesia, Tourette's syndrome, and
Parkinson's disease. Such compounds are also useful for inducing smoking cessation and are
25 neuroprotective.

SUMMARY OF THE INVENTION


The present invention encompasses spirofuopyridine aryl derivatives having activity
at nicotinic acetylcholine receptors ("nAChRs"). Spirofuopyridine aryl derivatives of the
30 invention are those in accord with formula I:

-2-




and pharmaceutically-acceptable salts thereof, wherein:

Ar is a moiety of formula II or III:



II



III,

A is O or S;

B is N at one or two occurrences and CR¹ at all other occurrences;

R¹ is independently at each occurrence hydrogen, -R², -C₂-C₆alkenyl, -C₂-C₆alkynyl, halogen, -CN, -NO₂, -NR³R⁴ or -OR⁵;

R² is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms, and 1 or 2 substituents selected from: -C₂-C₆alkenyl, -C₂-C₆alkynyl, -CN, -NR³R⁴, or -OR⁵;

R³ and R⁴ are independently at each occurrence hydrogen, R⁵, or in combination at any one occurrence of -NR³R⁴ are -(CH₂)_pJ(CH₂)_q wherein J is O, S, NH, NR⁵ or a bond;

R⁵ is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms;

p at each occurrence is 2, 3 or 4, and


q at each occurrence is 0, 1 or 2.

The invention also encompasses enantiomers, *in vivo*-hydrolysable precursors and pharmaceutically-acceptable salts of the spirofuropyridine aryl derivatives, pharmaceutical compositions and formulations containing them, methods of using them to treat diseases and conditions either alone or in combination with other therapeutically-active compounds or substances, processes and intermediates used to prepare them, uses of them as medicaments, uses of them in the manufacture of medicaments and uses of them for diagnostic and analytic purposes.

DETAILED DESCRIPTION OF THE INVENTION



In one aspect this invention comprises compounds that are potent ligands for nicotinic acetylcholine receptors (nAChRs).

Compounds of the invention are those in accord with formula I:



and pharmaceutically-acceptable salts thereof, wherein:

Ar is a moiety of formula II or III:



A is O or S;

B is N at one or two occurrences and CR¹ at all other occurrences;

R¹ is independently at each occurrence hydrogen, -R², -C₂-C₆alkenyl, -C₂-C₆alkynyl, halogen, -CN, -NO₂, -NR³R⁴ or -OR⁵;

R² is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms, and 1 or 2 substituents selected from: -C₂-C₆alkenyl, -C₂-C₆alkynyl, -CN, -NR³R⁴, or -OR⁵;

R³ and R⁴ are independently at each occurrence hydrogen, R⁵, or in combination at any one occurrence of -NR³R⁴ are -(CH₂)_pJ(CH₂)_q wherein J is O, S, NH, NR⁵ or a bond;

R⁵ is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms;


p at each occurrence is 2, 3 or 4, and

q at each occurrence is 0, 1 or 2.

Particular compounds of the invention are those in which B is N at one occurrence and 3 or 4 occurrences of R¹ are hydrogen. Other particular compounds of the invention are those in which B is N at two occurrences and 2 or 3 occurrences of R¹ are hydrogen

Other particular compounds of the invention are those in which B is N at one occurrence.

Other particular compounds of the invention are those in which Ar is formula IV:



Further particular compounds of the invention are those in which A is O.

Compounds of the invention include:

(2'R)-5'-(furo[3,2-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine};

(2'R)-5'-(furo[3,2-c]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine};

(2'R)-5'-(furo[2,3-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}; and

(2'R)-5'-(furo[2,3-c]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}.

The invention also relates to compounds according to formula I, their use in therapy and compositions containing them.

Another aspect of the invention relates to compounds according to formula I wherein one or more of the atoms is labelled with a radioisotope of the same element. In one embodiment this aspect of the invention the compound of formula I is labelled with tritium.

Another aspect the invention relates to the use of compounds according to formula I for the therapy of diseases mediated through the action of nicotinic acetylcholine receptors. One embodiment of this aspect of the invention relates to the use of compounds of formula I for the therapy of diseases mediated through the action of $\alpha 7$ nicotinic acetylcholine receptors. Another embodiment of this aspect of the invention relates to the use for therapy, wherein the condition or disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder. Another embodiment of this aspect of the invention relates to the use for therapy, wherein the disorder is anxiety, schizophrenia, or mania or manic depression. Another embodiment of this aspect of the invention relates to the use for therapy, wherein the disorder is Parkinson's disease,

Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses.

Another aspect of the invention relates to a use of a compound according to formula I in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial. One embodiment of this aspect of the invention relates to a use of a compound according to formula I in the manufacture of a medicament for the treatment or prophylaxis of neurological or psychotic disorders or intellectual impairment disorders. Another embodiment of this aspect of the invention relates to the use of a compound according to formula I in the manufacture of a medicament for the treatment or prophylaxis of jetlag, for inducing the cessation of smoking, for treating nicotine addiction including that resulting from exposure to products containing nicotine, craving, pain, and for ulcerative colitis.

Another aspect of the invention relates to a method of treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial which comprises administering a therapeutically effective amount of a compound according to formula I. One embodiment of this aspect of the invention relates to a method of treatment or prophylaxis of neurological or psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a compound according to formula I. Another embodiment of this aspect of the invention relates to a method of treatment, wherein the disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, or Attention Deficit Hyperactivity Disorder. Another embodiment of this aspect of the invention relates to a method of treatment, wherein the disorder is Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is loss of cholinergic synapses. Another embodiment of this aspect of the invention relates to a method of treatment, wherein the disorder is anxiety, schizophrenia or mania or manic depression. Yet another embodiment of this aspect of the invention relates to a method of treatment or prophylaxis of jetlag, treatment for inducing cessation of smoking, treatment for nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a compound according to formula I.

Another aspect of the invention relates to a pharmaceutical composition comprising a compound according to formula I, and a pharmaceutically-acceptable diluent or carrier.

Another embodiment of this aspect of the invention relates to a pharmaceutical composition for use in the treatment of prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial. Another embodiment of this aspect of the invention relates to a pharmaceutical composition for use in the treatment or prophylaxis of psychotic disorders or intellectual impairment disorders. Another embodiment of this aspect of the invention relates to a pharmaceutical composition for use in the treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, or mania or manic depression Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapse, jetlag, for inducing the cessation of smoking, for treating nicotine addiction including that resulting from exposure to products containing nicotine, craving, and for treating pain and ulcerative colitis. Yet another embodiment of this aspect of the invention relates to a pharmaceutical composition for treating or preventing a condition or disorder identified herein arising from dysfunction of nicotinic acetylcholine receptor neurotransmission in a mammal, preferably a human, comprising an amount of a compound of formula I, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, effective in treating or preventing such disorder or condition and an inert pharmaceutically-acceptable carrier.

For the uses and methods of treatment described herein the dosage administered will, of course, vary with the compound employed, the mode of administration and the treatment desired. However, in general, satisfactory results will be obtained when the compounds of the invention are administered at a daily dosage of from about 0.1 mg to about 20 mg/kg of animal body weight. Such doses may be given in divided doses 1 to 4 times a day or in sustained release form. For man, the total daily dose is in the range of from 5 mg to 1,400 mg, more preferably from 10 mg to 100 mg, and unit dosage forms suitable for oral administration comprise from 2 mg to 1,400 mg of the compound admixed with a solid or liquid pharmaceutical carrier or diluent.

The pharmaceutical compositions described herein will, of course, contain different quantities of compound depending on the compound employed. However, in general, compositions will provide from about 0.1 mg to about 20 mg per kg of animal body weight. Such doses may be given in divided doses 1 to 4 times a day or in sustained release form. For man, compositions will provide daily dose is in the range of from 5 mg to 1,400 mg, more

preferably from 10 mg to 100 mg, and will provide unit dosage forms suitable for oral administration from 2 mg to 1,400 mg of the compound admixed with a solid or liquid pharmaceutical carriers, lubricants or diluents.

The compounds of formula I, or an enantiomer thereof, and pharmaceutically-acceptable salts thereof, may be used on their own or in the form of appropriate medicinal preparations for enteral or parenteral administration. According to a further aspect of the invention, there is provided a pharmaceutical composition including preferably less than 80% and more preferably less than 50% by weight of a compound of the invention in admixture with an inert pharmaceutically-acceptable diluent or carrier.

Examples of diluents and carriers are:

- for tablets and dragees: lactose, starch, talc, stearic acid;
- for capsules: tartaric acid or lactose;
- for injectable solutions: water, alcohols, glycerine, vegetable oils;
- for suppositories: natural or hardened oils or waxes.

There is also provided a process for the preparation of such a pharmaceutical composition that comprises mixing the ingredients.

A further aspect of the invention is the use of a compound according to the invention, an enantiomer thereof or a pharmaceutically-acceptable salt thereof, in the manufacture of a medicament for the treatment or prophylaxis of one of the below mentioned diseases or conditions; and a method of treatment or prophylaxis of one of the above mentioned diseases or conditions, which comprises administering a therapeutically effective amount of a compound according to the invention, or an enantiomer thereof or a pharmaceutically-acceptable salt thereof, to a patient.

Compounds according to the invention are agonists of nicotinic acetylcholine receptors. While not being limited by theory, it is believed that agonists of the $\alpha 7$ nAChR (nicotinic acetylcholine receptor) subtype should be useful in the treatment or prophylaxis of psychotic disorders and intellectual impairment disorders, and have advantages over compounds that are or are also agonists of the $\alpha 4$ nAChR subtype. Therefore, compounds that are selective for the $\alpha 7$ nAChR subtype are preferred. The compounds of the invention are indicated as pharmaceuticals, in particular in the treatment or prophylaxis of psychotic disorders and intellectual impairment disorders. Examples of psychotic disorders include schizophrenia, mania and manic depression, and anxiety. Examples of intellectual impairment disorders include Alzheimer's disease, learning deficit, cognition deficit, attention deficit,

memory loss, and Attention Deficit Hyperactivity Disorder. The compounds of the invention may also be useful as analgesics in the treatment of pain (including chronic pain) and in the treatment or prophylaxis of Parkinson's disease, Huntington's disease, Tourette's syndrome, and neurodegenerative disorders in which there is loss of cholinergic synapses. The compounds may further be indicated for the treatment or prophylaxis of jetlag, for use in inducing the cessation of smoking, craving, and for the treatment or prophylaxis of nicotine addiction (including that resulting from exposure to products containing nicotine).

It is also believed that compounds according to the invention are useful in the treatment and prophylaxis of ulcerative colitis.

As used herein, the term "C₁₋₆ alkyl" refers to a straight-chained, branched, or cyclic C₁₋₆ alkyl group.

Radio-labelled forms

Another aspect of the invention relates to radio-labelled forms of the compounds of the invention. Such compounds that bind potently and selectively to a receptor are useful because sensitive and quantitative techniques are available for the detection of the radioactivity, which allow the interaction of a compound with its receptor to be detected and measured.

One method of discovering compounds that bind to a receptor is to perform a binding assay where the degree of displacement of a radio-labelled compound by another compound is measured. Thus, radio-labelled forms of compounds that potently bind receptors are useful to screen for novel medicinal compounds that bind to receptors. Such novel medicinal compounds may modulate the activity of those receptors by agonism, partial-agonism, or antagonism. Radio-labelled forms of compounds of the invention are therefore useful in a screen for the discovery of other compounds that bind to nicotinic receptors.

The ability of analogue compounds to bind to localized receptors within the body makes it possible to utilize such compounds for *in situ* imaging by PET, SPECT and similar imaging methods. PET imaging is accomplished with the aid of tracer compounds labelled with a positron-emitting isotope: Goodman, M. M. Clinical Positron Emission Tomography, Mosby Yearbook, 1992, K. F. Hubner et al., Chapter 14. For most biological targets, few isotopes are suitable. The carbon isotope, ¹¹C, has been used for PET, but its short half-life of 20.5 minutes limits its usefulness to compounds that can be synthesized and purified quickly, and to facilities that are proximate to a cyclotron where the precursor ¹¹C starting material is generated. Other more energetic isotopes have even shorter half-lives, ¹³N has a half-life of 10

minutes and ^{15}O has a half-life of two minutes. Nevertheless, PET studies have been carried out with these isotopes as described by Hubner, K. F., in Clinical Positron Emission Tomography, Mosby Year Book, 1992, K. F. Hubner, *et al.*, Chapter 2. [^{18}F]-labelled compounds have been used in PET studies, but their use is limited by the 110-minute half-life of the isotope. Most notably, [^{18}F]-fluorodeoxyglucose has been widely used in studies of glucose metabolism and localization of glucose uptake associated with brain activity. [^{18}F]-L-fluorodopa and other dopamine receptor analogues have also been used in mapping dopamine receptor distribution.


SPECT imaging employs isotope tracers that emit high energy photons (γ -emitters).

The range of useful isotopes is greater than for PET, but SPECT provides lower three-dimensional resolution. Nevertheless, SPECT is widely used to obtain clinically significant information about analogue binding, localization and clearance rates. A isotope used for SPECT imaging is ^{123}I , a γ -emitter with a 13.3 hour half life. Compounds labelled with ^{123}I can be shipped up to about 1000 miles from the manufacturing site, or the isotope itself can be transported for on-site synthesis. Eighty-five percent of the isotope's emissions are 159 KeV photons, which is readily measured by SPECT instrumentation currently in use.

Increasingly, the precise location and distribution of receptors in the brain and other tissues is of interest to clinical researchers, clinicians and diagnosticians. The distribution of nAChR's in the brains of individuals having disorders involving reduced cholinergic function such as Alzheimer's disease, cognitive or attention disorders, anxiety, depression, smoking cessation, neuroprotection, schizophrenia, analgesia, Tourette's syndrome, and Parkinson's disease is of growing interest as the molecular bases of these conditions is being discovered. The precise location and distribution of nAChRs in the brain and other tissues is also of importance in assessing the relevance of animal models of these conditions.

Methods of Preparation

Methods that may be used for the synthesis of compounds of formula I include the method outlined in Scheme 1. Unless otherwise noted Ar in Scheme 1 is as defined above for Formula 1.

**Scheme 1**

The compounds of formula I cross-coupling reaction of compounds of formula V and VI, wherein either X or Y is halogen or OSO_2CF_3 when, respectively, Y or X is an organometallic group. Suitable organometallic groups include boronic acid or boronic ester groups, $\text{B}(\text{OH})_2$, $\text{B}(\text{OAlk})_2$, or a trialkylstannyl group SnAlk_3 , wherein Alk is an alkyl group. The reaction is performed in the presence of a suitable organometallic catalyst and solvent. Suitable organometallic catalysts include palladium (0) complexes, for example tetrakis(triphenylphosphine)palladium(0) or a combination of tris(dibenzylideneacetone)dipalladium(0) and a suitable triarylphosphine or triarylsarsine ligand, for example triphenylphosphine, tri(o-tolyl)phosphine or triphenylarsine. Suitable solvents include inert ether solvents, for example 1,2-dimethoxyethane, tetrahydrofuran, or 1,4-dioxane, or alcohols, such as ethanol, or mixtures thereof. If the compound of formula V or VI is a boronic acid, the presence of a suitable base in addition to the other reagents is preferred. Suitable bases include sodium carbonate, cesium carbonate, and barium hydroxide. The reaction is carried out at a temperature of 0-120 °C, and preferably at a temperature of 60-120 °C.

Compounds of formula V wherein X is an organometallic group or compounds of formula VI, wherein Y is an organometallic group may be prepared from compounds of the corresponding formula wherein X or Y as appropriate is hydrogen, halogen, or OSO_2CF_3 by a suitable metallation or exchange procedure. The compounds wherein the organometallic group is $\text{B}(\text{OH})_2$ may be prepared from suitable aromatic compounds having hydrogen or halogen groups, by conversion to the corresponding aryllithium or arylmagnesium compounds followed by reaction with trialkylborate and subsequent hydrolysis of the resulting borate ester. Similarly, compounds wherein the organometallic group is a trialkylstannyl group may

be prepared from suitable aromatic compounds having hydrogen or halogen groups, by conversion to the corresponding aryllithium or arylmagnesium compounds followed by reaction with an appropriate trialkylstannyl halide. The formation of the aryllithium or arylmagnesium compound is performed in a suitable inert solvent, for example, tetrahydrofuran. Alternatively, the compounds wherein the organometallic group is $B(OH)_2$ may be prepared from suitable aromatic compounds having halogen or OSO_2CF_3 groups by reaction with bis(pinacolato)diboron and an organometallic catalyst, followed by hydrolysis of the resulting borate ester. Compounds wherein the said organometallic group is a trialkylstannyl group may be prepared from suitable aromatic compounds having halogen or OSO_2CF_3 groups by reaction with the appropriate bis(trialkyltin) in the presence of a suitable organometallic catalyst. The reaction is performed in a suitable inert solvent, for example tetrahydrofuran, and suitable organometallic catalyst include, for example tetrakis(triphenylphosphine). The reaction is performed at a temperature of about $0^\circ C$ to about $150^\circ C$, preferably about $20^\circ C$ to about $100^\circ C$. Typical procedures for effecting such conversions will be known to those of skill in the art.

Compounds of formula V or VI wherein X or Y as appropriate represents OSO_2CF_3 may be prepared from compounds of formula V or VI wherein X or Y represents OH by reaction with trifluoromethanesulfonic anhydride or other trifluoromethanesulfonylating agent in the presence of a base and a suitable solvent. Suitable bases include pyridine, and 2,6-di-*t*-butylpyridine. The reaction is preferably performed at a temperature of -78 to $120^\circ C$, and most preferably at a temperature of -78 to $0^\circ C$.

Compounds of formula V may be prepared by the methods described in international patent application publication number WO99/03859.

Compounds of formula VI may be prepared by methods described herein, are known in the literature, may be prepared by methods adapted from methods described herein or in the literature by a person skilled in the art of synthetic organic chemistry, or may be prepared by methods known or which will be apparent to a person skilled in the art of synthetic organic chemistry.

Radio-labelled forms of compounds of the invention are synthesized either by incorporating radio-labelled starting materials or, in the case of tritium, exchange of hydrogen for tritium by known methods. Known methods include (1) electrophilic halogenation, followed by reduction of the halogen in the presence of a tritium source, for example, by hydrogenation with tritium gas in the presence of a palladium catalyst, or (2) exchange of

hydrogen for tritium performed in the presence of tritium gas and a suitable organometallic (e.g. palladium) catalyst.

Where necessary, hydroxy, amino, or other reactive groups may be protected using a protecting group as described in the standard text "Protecting groups in Organic Synthesis", 3rd Edition (1999) by Greene and Wuts.

The above-described reactions, unless otherwise noted, are usually conducted at a pressure of about one to about three atmospheres, preferably at ambient pressure (about one atmosphere).

Unless otherwise stated, the above-described reactions are conducted under an inert atmosphere, preferably under a nitrogen atmosphere.

The compounds of the invention and intermediates may be isolated from their reaction mixtures by standard techniques.

Acid addition salts of the compounds of formula I which may be mentioned include salts of mineral acids, for example the hydrochloride and hydrobromide salts; and salts formed with organic acids such as formate, acetate, maleate, benzoate, tartrate, and fumarate salts. Acid addition salts of compounds of formula I may be formed by reacting the free base or a salt, enantiomer or protected derivative thereof, with one or more equivalents of the appropriate acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, e.g., water, dioxane, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed in vacuum or by freeze drying. The reaction may be a metathetical process or it may be carried out on an ion exchange resin.

The compounds of formula I exist in tautomeric or enantiomeric forms, all of which are included within the scope of the invention. The various optical isomers may be isolated by separation of a racemic mixture of the compounds using conventional techniques, e.g. fractional crystallisation, or chiral HPLC. Alternatively the individual enantiomers may be made by reaction of the appropriate optically active starting materials under reaction conditions that will not cause racemisation.

Pharmacology

The pharmacological activity of compounds of the invention may be measured using the tests set out below:

Test A - Assay for affinity at $\alpha 7$ nAChR subtype

[125 I]- α -Bungarotoxin (BTX) binding to rat hippocampal membranes. Rat hippocampi were homogenized in 20 volumes of cold homogenization buffer (HB: concentrations of constituents (mM): tris(hydroxymethyl)aminomethane 50; MgCl_2 1; NaCl 120; KCl 5; pH 7.4). The homogenate was centrifuged for 5 minutes at 1000 g, the supernatant was saved and the pellet re-extracted. The pooled supernatants were centrifuged for 20 minutes at 12000 g, washed, and resuspended in HB. Membranes (30–80 μg) were incubated with 5 nM [125 I]- α -BTX, 1 mg/mL BSA (bovine serum albumin), test drug, and either 2 mM CaCl_2 or 0.5 mM EGTA [ethylene glycol-bis(β -aminoethylether)] for 2 hours at 21 °C, and then filtered and washed 4 times over Whatman glass fibre filters (thickness C) using a Brandel cell harvester. Pretreating the filters for 3 hours with 1% (BSA/0.01% PEI (polyethyleneimine) in water was critical for low filter blanks (0.07% of total counts per minute). Nonspecific binding was described by 100 μM (–)-nicotine, and specific binding was typically 75%.

Test B - Assay for affinity to the $\alpha 4$ nAChR subtype

[^3H](–)-nicotine binding. Using a procedure modified from Martino-Barrows and Kellar (Mol Pharm (1987) 31:169-174), rat brain (cortex and hippocampus) was homogenized as in the [125 I]- α -BTX binding assay, centrifuged for 20 minutes at 12,000 x g, washed twice, and then resuspended in HB containing 100 μM diisopropyl fluorophosphate. After 20 minutes at 4 °C, membranes (approximately 0.5 mg) were incubated with 3 nM [^3H](–)-nicotine, test drug, 1 μM atropine, and either 2 mM CaCl_2 or 0.5 mM EGTA for 1 h at 4 °C, and then filtered over Whatman glass fiber filters (thickness C) (pretreated for 1 h with 0.5% PEI) using a Brandel cell harvester. Nonspecific binding was described by 100 μM carbachol, and specific binding was typically 84%.

Binding data analysis for Tests A and B

IC_{50} values and pseudo Hill coefficients (nH) were calculated using the non-linear curve-fitting program ALLFIT (DeLean A, Munson P J and Rodbard D (1977) Am. J. Physiol., 235:E97-E102). Saturation curves were fitted to a one site model, using the non-linear regression program ENZFITTER (Leatherbarrow, R.J. (1987)), yielding K_D values of 1.67 and 1.70 nM for the [125 I]- α -BTX and [^3H](–)-nicotine ligands respectively. K_i values were estimated using the general Cheng-Prusoff equation:

$$\text{K}_\text{i} = [\text{IC}_{50}] / ((2 + ([\text{ligand}] / [\text{K}_\text{D}])^n) / n - 1)$$

where a value of $n=1$ was used whenever $nH < 1.5$ and a value of $n=2$ was used when $nH \geq 1.5$. Samples were assayed in triplicate and were typically $\pm 5\%$. K_i values were determined using 6 or more drug concentrations. The compounds of the invention are compounds with binding affinities (K_i) of less than 1000 nM in either Test A or Test B, indicating that they are expected to have useful therapeutic activity.


The compounds of the invention have the advantage that they may be less toxic, be more efficacious, be longer acting, have a broader range of activity, be more potent, produce fewer side effects, are more easily absorbed or have other useful pharmacological properties.

Examples

Commercial reagents were used without further purification. *n*-Butyllithium was used as a solution in hexane. Mass spectra were recorded using an HPLC-MS system employing an HP-1100 HPLC and a Micromass LCZ Mass Spectrometer using APCI as the ionisation technique, an HPLC-MS system employing an HP-1100 HPLC and an HP-1100-series mass selective detector using APCI as the ionisation technique, or a GC-MS system employing an HP-6890 gas chromatograph and an HP-5973 mass selective detector employing electron impact ionisation, and are reported as m/z for the parent molecular ion. Room temperature refers to 20–25 °C. 5'-Bromospiro[1-azabicyclo[2.2.2]octane-3,2'-(3'H)-furo[2,3-b]pyridine] and other precursors were prepared as described in international patent application publication number WO 99/03859.

Preparation 1

(2'R)-5'-Trimethylstannyl-spiro[1-azabicyclo[2.2.2]octane-3,2'-(3'H)-furo[2,3-b]pyridine]




(2'R)-5'-Bromospiro[1-azabicyclo[2.2.2]octane-3,2'-(3'H)-furo[2,3-b]pyridine] (690 mg, 2.34 mmol), hexamethylditin (1.225 g, 0.27 mmol) and tetrakis(triphenylphosphine)palladium (0) (266 mg, 0.027 mmol) were mixed with 10 mL of toluene and sealed under nitrogen. The mixture was stirred and heated at 120 °C under nitrogen for 4 h. The mixture was then allowed to cool and filtered through diatomaceous earth. The filtrate was diluted with chloroform, washed with saturated sodium bicarbonate, dried through $MgSO_4$, filtered, and the solvent was evaporated. Purification by flash

chromatography using a gradient of ammoniated methanol in chloroform gave the title compound as a solid (780 mg); m/e 377 379 381 (MH⁺).

Example 1

5 (2'R)-5'-(Furo[3,2-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}



(a) Furo[3,2-b]pyridin-3-yl trifluoromethanesulfonate

3-Hydroxy-picolinic acid (15.6 g, 42.1 mmol), ethanol (360 mL), benzene (100 mL) and 98% sulfuric acid (6 mL) were heated under reflux for 40 h. After evaporation of the ethanol and benzene, the residue was dissolved in water, neutralized with sodium bicarbonate, and extracted with chloroform. The organic layer was dried (MgSO₄), filtered, and then the solvent was evaporated to give ethyl 3-hydroxy-picolinate as a brown oily residue (11.0 g).

A mixture of ethyl 3-hydroxy-picolinate (11.0 g, 65.8 mmol), ethyl bromoacetate (12.1 g, 72.4 mmol) and anhydrous potassium carbonate (11.8 g, 85.5 mmol) in acetone (120 mL) was heated under reflux for 15 h. After cooling, the inorganic material separated by filtration. The filtrate was dissolved in chloroform, washed with water, then brine and dried (MgSO₄), filtered, and the solvent was evaporated to give a brown oily residue. The residue was purified by flash chromatography using a gradient of ammoniated methanol in chloroform to give ethyl 2-(2-ethoxycarbonyl-3-pyridyloxy) acetate (13.7 g) as a yellow oil (13.7 g).

Ethyl 2-(2-ethoxycarbonyl-3-pyridyloxy) acetate (13.6 g, 54.0 mmol) and sodium ethoxide (8.08 g, 118.8 mmol) in toluene (200 mL) were heated under reflux for 18 h. After cooling, a precipitate was collected by filtration, dissolved in the minimum amount of hot water (about 300 mL), and acidified with acetic acid (6 mL). The resulting precipitate was filtered, and dried *in vacuo* to give ethyl 3-hydroxyfuro[3,2-b]pyridine-2-carboxylate as a solid (7.0 g).

Ethyl 3-hydroxyfuro[3,2-b]pyridine-2-carboxylate (6.90 g, 25.8 mmol) was dissolved in 10% hydrochloric acid (50 mL), and heated under reflux for 3 h. Evaporation of the hydrochloric acid solution gave furo[3,2-b]pyridin-3(2H)-one hydrochloride (9.0 g). A portion of the furo[3,2-b]pyridin-3(2H)-one hydrochloride was converted to its free base by

treatment with saturated sodium bicarbonate and extraction with chloroform in preparation for the following step.


N,N-Diisopropylethylamine (1.08 g, 8.34 mmol) was added slowly to a solution of furo[3,2-b]pyridin-3(2H)-one (1.40 g, 7.25 mmol) in dry dichloromethane (60 mL) under nitrogen at -10°C . Then trifluoromethanesulfonic anhydride (2.45 g, 8.70 mmol) was added slowly. The mixture was warmed to RT, and stirred overnight. The reaction was quenched with water. The organic layer was washed with water and brine, dried (MgSO_4), and then the solvent was evaporated to give a brown oily residue, which was purified by flash chromatography using chloroform to give the sub-title compound (980 mg) as a light-brown oil.

(b) (2'R)-5'-(Furo[3,2-b]pyridine-3-yl)spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine]

(2'R)-5'-Trimethylstannyl-spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine] (200 mg, 0.53 mmol), furo[3,2-b]pyridine-3-triflate (184 mg, 0.69 mmol), Pd_2dba_3 (25 mg, 0.027 mmol) and triphenylphosphine (14 mg, 0.053 mmol) were mixed in 2 mL of DMF. The reaction was heated to 100°C for six hours. After cooling, the mixture was poured into a solution of brine, and extracted with chloroform. The organic layer was dried through MgSO_4 , and the solvent was evaporated. The residue was purified by flash chromatography using a gradient of ammoniated methanol in chloroform to give a solid. The solid was further purified by reverse phase HPLC on a Waters Novapak-HR C_{18} Column using a gradient of 5 - 45% acetonitrile/water as the eluant (each solvent containing 0.1% trifluoroacetic acid as a buffer). The product-containing fractions were evaporated. The residue was neutralized with NaHCO_3 , extracted with chloroform, and the solvent was evaporated to give the title compound (33 mg) as a white solid; m/e 334.3 (MH^+).

Example 2

(2'R)-5'-(Furo[3,2-c]pyridine-3-yl)spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine]



(a) Furo[3,2-c]pyridin-3-yl trifluoromethanesulfonate

To a solution of 4-chloronicotinic acid (7.03 g, 44.6 mmol) in 100 mL of hexamethylphosphoric triamide (HMPA) was added a solution of 5.5 N sodium hydroxide (13 mL) at 0 °C. Then iodoethane (27.8 g, 179 mmol) was added into the reaction mixture at 0 °C over 1 h, and stirring was continued for a further 1 h at room temperature. The mixture was diluted with 250 mL of water, and then extracted three times with ether. The ethereal extracts were combined, washed three times with water, then dried (MgSO₄), filtered, and the solvent was evaporated to give ethyl 4-chloronicotinate (6.21 g) as a brown oily residue, which was used for the next step without further purification.

Ethyl glycolate (7.26 g, 59.7 mmol) was added slowly into a suspension of sodium hydride (2.95 g of 60% dispersion in mineral oil, 73.8 mmol) in 85 mL of 1,2-dimethoxyethane (DME) at 0 °C, and the mixture was stirred for another 30 min. A solution of ethyl 4-chloronicotinate (6.20 g, 33.4 mmol) in 20 mL of DME was added slowly into the reaction mixture at room temperature. The mixture was heated to 70 °C and maintained at that temperature overnight. The solvent was evaporated and the residue was dissolved in 100 mL of water, and washed with hexane 3 times. The pH of the water solution was adjusted to about 5 using acetic acid to pH 5, and a yellow precipitate was formed. The yellow precipitate was filtered, washed with a small amount of water (20 mL x 3), and dried overnight *in vacuo* to give ethyl 3-hydroxyfuro[3,2-c]pyridine-2-carboxylate free base as a yellow solid (5.79 g) which was used for the next step without further purification.


Ethyl 3-hydroxyfuro[3,2-c]pyridine-2-carboxylate (5.79 g, 27.9 mmol) was dissolved in 10% hydrochloric acid (50 mL), and heated under reflux for 40 h. The reaction mixture was evaporated to dryness. The residue was suspended in saturated sodium bicarbonate solution and extracted with chloroform 3 times. The chloroform extracts were combined, dried (MgSO₄), filtered, and evaporated to give furo[3,2-c]pyridin-3(2H)-one (570 mg) as a light-brown oil. N,N-Diisopropylethylamine (616 mg, 4.77 mmol) was added slowly to a solution of furo[3,2-c]pyridin-3(2H)-one (560 mg, 4.14 mmol) in dry methylene chloride (25 mL) under nitrogen at -10 °C. Then trifluoromethanesulfonic anhydride (1.40 g, 4.97 mmol) was added slowly. The mixture was allowed to warm to RT, and stirred for another 1 g. The reaction was quenched with water. The organic layer was washed with water, dried (MgSO₄), filtered, and then the solvent was evaporated to give the sub-title compound as a brown oily residue (1.10 g), which was used without further purification for the next step.

(b) (2'R)-5'-(Furo[3,2-c]pyridine-3-yl)spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine]

Prepared by a method analogous to that described for the preparation of Example 1 from (2'R)-5'-trimethylstannyl-spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine] and furo[3,2-c]pyridin-3-yl trifluoromethanesulfonate. After reverse phase chromatography, the title compound was obtained as the trifluoroacetate salt which was a colourless solid; m/e 334 (MH⁺).

Example 3

(2'R)-5'-(Furo[2,3-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}



(a) Furo[2,3-b]pyridin-3-yl trifluoromethanesulfonate

2-Chloronicotinic acid (5.90 g, 37.5 mmol), ethanol (120 mL), toluene (35 mL) and 98% sulfuric acid (2 mL) were heated together under reflux for 48 hours. After evaporation of the ethanol and toluene, the residue was dissolved in water, neutralized with sodium bicarbonate, and extracted with chloroform. The organic layer was dried through MgSO₄, and then the solvent was evaporated to give a dark-brown oil-like residue, which was purified by flash chromatography using a gradient of 10 - 20% hexane/ethyl acetate as the eluant to give ethyl 2-chloronicotinate (4.60 g) as a light-brown oil.

Ethyl glycolate (7.00 g, 67.3 mmol) was added slowly into a suspension of sodium hydride (2.80 g of 60% dispersion in mineral oil, 70.0 mmol) in 60 mL of 1,2-dimethoxyethane (DME) at 0 °C, and the mixture was then stirred for another 30 min. A solution of ethyl 2-chloronicotinate (4.60 g, 24.8 mmol) in 14 mL of DME was added into the reaction mixture slowly at room temperature. Then the mixture was heated at 70 °C overnight. After evaporation of the solvent, the residue was dissolved in 90 mL of water, washed three times with hexane, acidified with acetic acid, and extracted three times with chloroform. The combined chloroform layers were dried (MgSO₄), and evaporated to give a yellow residue. The residue was crystallized from 20 mL of ether to give ethyl 3-hydroxyfuro[2,3-b]pyridine-2-carboxylate as a pale-yellow solid (4.30 g).

Ethyl 3-hydroxyfuro[2,3-b]pyridine-2-carboxylate (3.0 g, 11.23 mmol) was heated under reflux in 10 % hydrochloric acid (50 mL) for 3 h. Evaporation of the hydrochloric acid

solution gave furo[2,3-b]pyridin-3(2H)-one as the hydrochloride salt (1.73 g). A portion of the furo[2,3-b]pyridin-3(2H)-one hydrochloride was converted to its free base by treated with saturated sodium bicarbonate and extracted with chloroform in preparation for the following step.


5 N,N-Diisopropylethylamine (429 mg, 3.32 mmol) was added slowly to a solution of furo[2,3-b]pyridin-3(2H)-one (390 mg, 3.89 mmol) in dry dichloromethane under nitrogen at -10 °C. Then, trifluoromethanesulfonic anhydride (978 mg, 3.47 mmol) was added slowly. The mixture was allowed to warm to RT, and stirred for another 2 h. The reaction was quenched with water. The organic layer was washed with water, and then dried (MgSO₄),
10 filtered, and then the solvent was evaporated to give a brown oily residue. Purification by flash chromatography using chloroform to give the sub-title compound as a light-brown oil (576 mg).

(b) (2'R)-5'-(Furo[2,3-b]pyridine-3-yl)spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine]

15 Prepared by a method analogous to that described for the preparation of Example 1 from (2'R)-5'-trimethylstannyl-spiro[1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine] and furo[2,3-b]pyridin-3-yl trifluoromethanesulfonate. The title compound was obtained as a colourless solid; m/e 334 (MH⁺).

20 Example 4


(2'R)-5'-(Furo[2,3-c]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}



The title compound is prepared by a synthesis analogous to that described in the
25 synthesis of Example 1 starting from 3-hydroxypyridine-4-carboxylic acid [Di Marco *et al.*,
Eur. J. Inorg. Chem., 2002, (10), 2648-2655].


CLAIMS

1. A compound of formula I:




- 5 and pharmaceutically-acceptable salts thereof, wherein:

Ar is a moiety of formula II or III:



II



III,

wherein,

- 10 A is O or S;

B is N at one or two occurrences and CR¹ at all other occurrences;

R¹ is independently at each occurrence hydrogen, -R², -C₂-C₆alkenyl, -C₂-C₆alkynyl, halogen, -CN, -NO₂, -NR³R⁴ or -OR⁵;

- 15 R² is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms, and 1 or 2 substituents selected from: C₂-C₆alkenyl, C₂-C₆alkynyl, -CN, -NR³R⁴, or -OR⁵;

R³ and R⁴ are independently at each occurrence hydrogen, R⁵, or in combination at any one occurrence of -NR³R⁴ are -(CH₂)_pJ(CH₂)_q wherein J is O, S, NH, NR⁵ or a bond;

- 20 R⁵ is an unsubstituted straight-chained, branched, or cyclic C₁-C₆alkyl group, or a straight-chained, branched, or cyclic C₁-C₆alkyl group substituted with 1, 2, 3, 4 or 5 halogen atoms;

p at each occurrence is 2, 3, or 4;

q at each occurrence is 0, 1, or 2.


25

2. A compound according to Claim 1 or a pharmaceutically-acceptable salt thereof,

selected from compounds wherein B is N at one occurrence and 2 or 3 occurrences of R¹ are hydrogen, or compounds wherein B is N at two occurrences and 3 or 4 occurrences of R¹ are hydrogen.

3. A compound according to Claim 1 wherein B is N at one occurrence.

4. A compound according to Claim 1 or a pharmaceutically-acceptable salt thereof, wherein Ar is formula IV:



5. A compound according to Claim 1 or a pharmaceutically-acceptable salt thereof, wherein A is O.

6. A compound according to Claim 1, selected from the group consisting of:

(2'R)-5'-(furo[3,2-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine};

(2'R)-5'-(furo[3,2-c]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine};

(2'R)-5'-(furo[2,3-b]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}; and

(2'R)-5'-(furo[2,3-c]pyridine-3-yl)spiro{1-azabicyclo[2.2.2]octane-3,2'(3'H)-furo[2,3-b]pyridine}.

7. A pharmaceutical composition comprising a compound according to Claim 1, and a pharmaceutically-acceptable diluent or carrier.

8. The pharmaceutical composition according to Claim 7, for use in the treatment of prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial.

9. The pharmaceutical composition according to Claim 7, for use in the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

5 10. The pharmaceutical composition according to Claim 7, for use in the treatment or prophylaxis of Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder, anxiety, schizophrenia, or mania or manic depression Parkinson's disease, Huntington's disease, Tourette's syndrome, neurodegenerative disorders in which there is loss of cholinergic synapse, jetlag, smoking,
10 addiction to tobacco, nicotine addiction including that resulting from exposure to products containing nicotine, craving, pain, and for ulcerative colitis.

11. Use of a compound according to Claim 1, in the manufacture of a medicament for the treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$
15 nicotinic receptor is beneficial.

12. Use of a compound according to Claim 1, in the manufacture of a medicament for the treatment or prophylaxis of neurological disorders, psychotic disorders or intellectual impairment disorders.

20

13. The use according to Claim 12, wherein the condition or disorder is Alzheimer's disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder.

25 14. The use according to Claim 12, wherein the disorder is anxiety, schizophrenia, or mania or manic depression.

15. The use as claimed in claim 12, wherein the disorder is Parkinson's disease, Huntington's disease, Tourette's syndrome, or neurodegenerative disorders in which there is
30 loss of cholinergic synapses.

16. Use of a compound according to Claim 1 in the manufacture of a medicament for the treatment or prophylaxis of jetlag, cessation of smoking, addiction to tobacco, nicotine addiction including that resulting from exposure to products containing nicotine, craving, pain, and for ulcerative colitis.

5

17. A method of treatment or prophylaxis of human diseases or conditions in which activation of the $\alpha 7$ nicotinic receptor is beneficial which comprises administering a therapeutically effective amount of a compound according to Claim 1.

10 18. A method of treatment or prophylaxis of psychotic disorders or intellectual impairment disorders, which comprises administering a therapeutically effective amount of a compound according to Claim 1.

19. The method according to Claim 18, wherein said psychotic disorder is Alzheimer's
15 disease, learning deficit, cognition deficit, attention deficit, memory loss, Attention Deficit Hyperactivity Disorder Parkinson's disease, Huntington's disease, Tourette's syndrome, a neurodegenerative disorder in which there is loss of cholinergic synapses anxiety, schizophrenia or mania or manic depression.

20 20. A method of treatment or prophylaxis of jetlag, cessation of smoking, nicotine addiction, craving, pain, and for ulcerative colitis, which comprises administering a therapeutically effective amount of a compound according to Claim 1.

21. A compound according to Claim 1, wherein one or more of the atoms is a radioisotope
25 of the element.

22. A compound according to Claim 21, wherein the radioisotope is tritium.

23. The use of a compound according to Claim 21, in a screen for the discovery of novel
30 medicinal compounds which bind to and modulate the activity, *via* agonism, partial agonism, or antagonism, of the $\alpha 7$ nicotinic acetylcholine receptor.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/GB2004/004484

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D491/04 A61K31/438 A61P25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00/42044 A (ASTRAZENECA AB; LOCH, JAMES, III; MULLEN, GEORGE; PHILLIPS, EIFION) 20 July 2000 (2000-07-20) claim 1; page 18ff	1-23
A	WO 02/096912 A (ASTRAZENECA AB; PHILLIPS, EIFION) 5 December 2002 (2002-12-05) claim 1; page 2	1-23
P, A	WO 03/087102 A (ASTRAZENECA AB; CHANG, HUI-FANG; LI, YAN; PHILLIPS, EIFION) 23 October 2003 (2003-10-23) p. 3ff; claim 1	1-23
P, A	WO 03/087103 A (ASTRAZENECA AB; CHANG, HUI-FANG; LI, YAN; PHILLIPS, EIFION) 23 October 2003 (2003-10-23) p.3ff; claim 1	1-23

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *A* document member of the same patent family

Date of the actual completion of the international search

7 April 2005

Date of mailing of the international search report

14/04/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Wolf, C

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box II.1

Although claims 17-20 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition. Although claim 23 is directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Continuation of Box II.1

Claims Nos.: claims 17-20,23

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

Rule 39.1(iv) PCT - Diagnostic method practised on the human or animal body

INTERNATIONAL SEARCH REPORT

International application No.
PCT/GB2004/004484

Box II Observations where certain claims were found unsearchable (Continuation of Item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: claims 17-20, 23
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB2004/004484

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0042044	A	20-07-2000	AT 240960 T 15-06-2003
			AU 775433 B2 29-07-2004
			AU 2334300 A 01-08-2000
			BR 9916906 A 30-10-2001
			CA 2359990 A1 20-07-2000
			CN 1132837 C 31-12-2003
			CZ 20012554 A3 16-01-2002
			DE 69908185 D1 26-06-2003
			DE 69908185 T2 03-06-2004
			DK 1147114 T3 15-09-2003
			EE 200100370 A 15-10-2002
			EP 1147114 A1 24-10-2001
			ES 2200590 T3 01-03-2004
			HK 1040517 A1 05-09-2003
			HU 0200513 A2 29-06-2002
			ID 30034 A 01-11-2001
			JP 2002534525 T 15-10-2002
			NO 20013478 A 14-09-2001
			NZ 512733 A 29-08-2003
			PT 1147114 T 31-10-2003
			RU 2233282 C2 27-07-2004
			WO 0042044 A1 20-07-2000
			SI 1147114 T1 31-12-2003
			SK 9932001 A3 04-04-2002
			TR 200102042 T2 21-05-2002
			US 2003149065 A1 07-08-2003
			ZA 200105527 A 04-10-2002
WO 02096912	A	05-12-2002	BR 0210075 A 17-08-2004
			CA 2455341 A1 05-12-2002
			CN 1512995 A 14-07-2004
			EP 1397366 A1 17-03-2004
			JP 2004532877 T 28-10-2004
			MX PA03010996 A 27-02-2004
			WO 02096912 A1 05-12-2002
WO 03087102	A	23-10-2003	AU 2003225456 A1 27-10-2003
			BR 0309343 A 15-02-2005
			CA 2482311 A1 23-10-2003
			EP 1499618 A1 26-01-2005
			WO 03087102 A1 23-10-2003
WO 03087103	A	23-10-2003	AU 2003224545 A1 27-10-2003
			BR 0309342 A 15-02-2005
			CA 2482312 A1 23-10-2003
			EP 1499615 A1 26-01-2005
			WO 03087103 A1 23-10-2003